AD	)				

Award Number: W81XWH-10-1-0715

TITLE: Magnetic Resonance Characterization of Axonal Response to Spinal Cord

Injury

PRINCIPAL INVESTIGATOR: Alan Tessler, Ph.D.

CONTRACTING ORGANIZATION: Drexel University

Philadelphia, PA 19104

REPORT DATE: October 2011

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

**Distribution Unlimited** 

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

data needed, and completing this burden to Department of	and reviewing this collection of i Defense, Washington Headquar	nformation. Send comments reg ters Services, Directorate for Info	arding this burden estimate or ar rmation Operations and Reports	ny other aspect of this of (0704-0188), 1215 Jeff	ching existing data sources, gathering and maintaining the ollection of information, including suggestions for reducing erson Davis Highway, Suite 1204, Arlington, VA 22202-		
4302. Respondents should be valid OMB control number. <b>P</b>	e aware that notwithstanding and LEASE DO NOT RETURN YOU	y other provision of law, no perso IR FORM TO THE ABOVE ADD	n shall be subject to any penalty RESS.	for failing to comply wit	h a collection of information if it does not display a currently		
1. REPORT DATE		2. REPORT TYPE			DATES COVERED		
October 2011		Annual			September 2010 – 26 September 2011		
4. TITLE AND SUBTIT	「LE			5a.	CONTRACT NUMBER		
Magnetic Resona	nce Characterizatio	n of Axonal Respon	ise to Spinal Cord Ir	ijui y	5b. GRANT NUMBER W81XWH-10-1-0715		
				5c.	PROGRAM ELEMENT NUMBER		
0 AUTUOD(0)					DD 0 ISOT NUMBER		
6. AUTHOR(S)				50.	PROJECT NUMBER		
Alexa Terralesa Dia I	_			Fo	TASK NUMBER		
Alan Tessler, Ph.	J.			5e.	TASK NUMBER		
				Ef.	WORK UNIT NUMBER		
E-Mail: wehrli@mail.r	nod upopp odu			31.	WORK ONLY NOWIDER		
	GANIZATION NAME(S)	AND ADDDESS(ES)		0 1	PERFORMING ORGANIZATION REPORT		
7. FERFORMING OR	SANIZATION NAME(S)	AND ADDRESS(ES)			NUMBER		
Drexel University							
Philadelphia, PA	19104						
9. SPONSORING / MO	ONITORING AGENCY N	NAME(S) AND ADDRES	S(ES)	10.	SPONSOR/MONITOR'S ACRONYM(S)		
	I Research and Ma		0(20)	101			
Fort Detrick, Mary							
	= 0= 00 .=			11.	SPONSOR/MONITOR'S REPORT		
					NUMBER(S)		
12. DISTRIBUTION / /	VAILABILITY STATE	/ENT					
	ic Release; Distribu						
	, , , , , , , , , , , , , , , , , , , ,						
13. SUPPLEMENTAR	V NOTES						
13. SOFFLEWENTAN	INOTES						
14. ABSTRACT							
None provided							
None provided							
45 0110 1505 5555							
15. SUBJECT TERMS	j						
None provided.							
			1	1			
16. SECURITY CLAS	SIFICATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER	19a. NAME OF RESPONSIBLE PERSON		
	T		OF ADSTRACT	OF PAGES	USAMRMC		
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (include area		
U	U	U	UU	5	code)		

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

#### INTRODUCTION

Spinal cord injuries (SCI) produce direct mechanical disruption and subsequent severe degeneration of axons; these processes cause the associated neurologic deficits. Histological studies of fixed tissue in animal models of SCI have described axonal loss and demyelination. Research at the U. Penn site brings novel magnetic resonance methodology to bear with the objective of obtaining quantitative information on axonal degeneration and myelin loss following SCI in a mouse model by pursuing the following specific aims per the work statement:

- 1. We will perform q-space MR imaging (QSI) and simulations of QSI to quantify axonal architecture in healthy and injured mouse spinal cords.
- 2. We will quantify myelin content with three quantitative MRI techniques in healthy and injured mouse spinal cords and compare the results with histology.

# Specific Aim 1:

Normal and injured mice have been prepared for QSI analysis. A small pilot set of spinal cord specimens was used to refine the imaging approaches (normal & injured). Technical details have been worked out regarding injury placement, tissue collection and marking to enable reliable identification of the lesion site and rostral/caudal orientation of the tissue specimens. We have generated healthy (n=4) and injured spinal cord tissue (2-day, 3-week and 3-month sacrifice, n=4 per time point).

Upgrading the Bruker NMR/MRI system has delayed QSI experiments. Hardware modifications were needed to connect our previous custom gradient coil to the new system. The gradient coil also had to be optimized and recalibrated for the new system. As the old QSI pulse sequence program does not run on the new system, a new QSI pulse sequence program is currently under development.

Once the new pulse sequence program has been tested, QSI experiments will be performed and followed by histologic analysis and QSI simulations.

There has been significant progress toward translation of the QSI methodology to the clinic. Toward this goal a pulse sequence was designed and implemented for generating a series of images as a function of q (the spatial wave vector) at 1.5T on a clinical imager. Using this pulse sequence on fixed pig spinal cords, we have collected preliminary data to investigate the feasibility of using our previously published QSI methods on a clinical scanner.

#### Specific Aim 2:

Significant progress has been made towards 3D ultra-short echo-time (UTE) MRI of myelin. First, we succeeded in isolating bovine myelin and demonstrated that the spectroscopic and imaging characteristics of the hydrated myelin were identical to those obtained in situ in rat spinal cord. In the course of these experiments the MR signal of myelin was studied extensively with proton, carbon and phosphorus NMR spectroscopy. The results of this pilot study indicate that UTE MRI may have potential for directly imaging myelin. We

demonstrated the feasibility of a 3D dual-echo subtraction UTE sequence with adiabatic inversion long- $T_2$  suppression to directly image myelin in a freshly excised rat spinal cord (Figure 1). Lastly, we demonstrated a quantitative relationship between image-derived signal intensities and actual myelin concentration.




Figure 1. Sample 3D UTE images from rat thoracic spinal cords averaged over five central slices. Images obtained for a) TE=10 ms, b) TE=1,200 ms, and c) magnitude difference (maximum intensity range decreased by a factor of two to highlight myelin signal). D) Intensity profiles across the three images (delineated as red, green and yellow lines in panels a, b and c, respectively) to show relative white matter (WM), grey matter (GM) and background intensity. The dashed blue line represents the average noise level. WM and GM are indicated in panel a, and arrows highlight residual surface water in panel b.

Similar to Specific Aim 1, progress toward detecting myelin with IHMT and MR relaxometry has been hampered by the disruption in imaging capabilities caused by the upgrade of the Bruker Instruments micro-imaging system. However, as of the time of this report (10-15-11) software and hardware upgrades are complete and work should resume in November 2011.

#### KEY RESEARCH ACCOMPLISHMENTS

- Demonstrated feasibility of direct imaging of neural myelin as a new metric for the evaluation of SCI.
- Magnetic resonance characterization and feasibility demonstration of MR imaging of myelin in situ has been presented at the International Society for Magnetic Resonance in Medicine (ISMRM) Annual Meeting in Montreal, Canada (May 2011) and American Society for Neuroradiology Annual Meeting in Seattle (June 2011). See citations below.
- Submitted a manuscript on myelin MRI for publication to the Proceedings of the National Academy of Science.
- Generated the model injury mouse spinal cords.
- Tested the system upgrade of the Bruker Instruments micro-imaging system and interfaced custom-built gradients for high-resolution q-space imaging.
- Demonstrated the feasibility of translation of the q-space imaging technique in porcine model of the spinal cord on a 1.5T clinical imager and an abstract is being submitted for presentation at the ISMRM Annual Meeting in Melbourne, Australia in 2012.

## OUTCOMES

The new myelin imaging technique has shown potential for quantitative assessment of myelin content in the CNS of the rat spinal cord.

### CONCLUSION

While the project is slightly delayed, the progress made during the first year of the project gives the investigators confidence that the project will be completed in a timely fashion.

#### REFERENCES

- 1. Wilhelm MJ, Ong HH, Wehrli SL Tsai P-H, Hackney DB, Wehrli FW. Prospects for quantitative imaging of myelin with dual-echo short inversion time 3D UTE MRI. Proc. Intl. Soc. Mag. Reson. Med. 19 (2011). Montreal, Quebec, Canada. P. 2640.
- 2. Wilhelm MJ, Ong HH, Wehrli SL, Tsai P-H, Wright AC, Hackney DB, Wehrli FW. Prospects for quantitative imaging of myelin with ultra-short TE 3D radial MRI. 49th ASNR, Seattle, WA, June 4-9, (2011).